

TN:2251135



Journal Title: American Mathematical Society translations, series 2, vol. 17.

Volume:

Issue: 17

Month/Year: 1961

Pages: 369-373

Article Author: Kolmogorov, A.N.

Article Title: On the representation of continuous functions of several variables by superpositions of continuous functions of a smaller number of variables

Cited In:

Notes:5/31/2018 7:35:24 AM

(sokolov@uchicago.edu)

<https://www.ams.org/books/trans2/017/>

6/19/2018 11:06:06 AM (lashanda)

@Mansueto

6/19/2018 11:09:18 AM (System) IDS

Logic::LogicRule::FlagForMansueto1::Processed

Print Date:6/19/2018 11:24 AM

Call #: QA1.A545 c.1 no.11-20

Location: ASR / SciASR

Barcode:11315227



ODYSSEY REQUEST

Vadim Sokolov
sokolov@uchicago.edu

Sent S.H. 6/19/18

ODYSSEY REQUEST

Notice: This material may be protected
by copyright law (Title 17 US Code)



ILLiad TN: 2251135

Copyright © 1961 by the American Mathematical Society
 Printed in the United States of America

All Rights Reserved
 No portion of this book may be
 reproduced without the written
 permission of the publisher

Library of Congress Catalog Card Number A51-5559

PHOTOLITHOPRINTED BY CUSHING - MALLOY, INC.
 ANN ARBOR, MICHIGAN, UNITED STATES OF AMERICA

TABLE OF CONTENTS

	Page
Plotkin, B. I. On some criteria of locally nilpotent groups. [Плоткин, Б. И. О некоторых признаках локально нильпотентных групп.] <i>Uspehi Mat. Nauk (N.S.)</i> 9 (1954), no. 3(61), 181-186.....	1
Plotkin, B. I. Radical groups. [Плоткин, Б. И. Радикальные группы.] <i>Mat. Sb. N.S.</i> 37(79) (1955), 507-526.....	9
Plotkin, B. I. Generalized solvable and generalized nilpotent groups. [Плоткин, Б. И. Обобщенные разрешимые и обобщенные нильпотентные группы.] <i>Uspehi Mat. Nauk (N.S.)</i> 13 (1958), no. 4(82), 89-172.....	29
Černikov, S. N. Groups with systems of complemented subgroups. [Черников, С. Н. Группы с системами дополняемых подгрупп.] <i>Mat. Sb. N.S.</i> 35(77) (1954), 93-128.....	117
Černikova, N. V. Groups with complemented subgroups. [Черникова, Н. В. Группы с дополняемыми подгруппами.] <i>Mat. Sb. N.S.</i> 39(81) (1956), 273-292.....	153
Mal'cev, A. I. Free topological algebras. [Мальцев, А. И. Свободные топологические алгебры.] <i>Izv. Akad. Nauk SSSR. Ser. Mat.</i> 21 (1957), 171-198.....	173
Rimskii-Korsakov, B. S. A version of the construction of a theory of generalized gamma-functions based on the Laplace transform. [Римский-Корсаков, Б. С. Вариант построения теории обобщенных гамма-функций на базе преобразования Лапласа.] <i>Moskov. Oblast. Pedagog. Inst. Uč. Zap.</i> 57 (1957), 121-141.....	201
Vilenkin, N. Ya. Supplement to "Theory of Orthogonal Series" (Russian translation of the book by S. Kaczmarz and H. Steinhaus, translated with a reviewing article by R. S. Guter and P. L. Ulyanov, edited with a supplement by N. Ya. Vilenkin.) [Виленкин, Н. Я. Дополнения Теория ортогональных рядов.] <i>Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow</i> , 1958.....	219
Trohimčuk, Yu. Yu. A theorem of H. Bohr and its generalizations. [Трохимчук, Ю. Ю. Теорема Г. Бора и ее обобщения.] <i>Mat. Sb. N.S.</i> 45(87) (1958), 233-260.....	251
Kolmogorov, A. N.; and Tihomirov, V. M. ϵ -entropy and ϵ -capacity of sets in functional spaces. [Колмогоров, А. Н.; и Тихомиров, В. М. ϵ -энтропия и ϵ -емкость множеств в функциональных пространствах.] <i>Uspehi Mat. (N.S.)</i> 14 (1959), no. 2(86), 3-86.....	277
Vituškin, A. G. The absolute ϵ -entropy of metric spaces. [Витушкин, А. Г. Абсолютная ϵ -энтропия метрических пространств.] <i>Dokl. Akad. Nauk SSSR (N.S.)</i> 117 (1957), 745-747.....	365
Kolmogorov, A. N. On the representation of continuous functions of sev-	

**ON THE REPRESENTATION OF CONTINUOUS FUNCTIONS
OF SEVERAL VARIABLES BY SUPERPOSITIONS OF CONTINUOUS FUNCTIONS
OF A SMALLER NUMBER OF VARIABLES**

A. N. KOLMOGOROV

From Theorem 3, stated below, there follows a rather unexpected consequence: every continuous function of an arbitrarily large number of variables can be represented in the form of a finite superposition of continuous functions of not more than three variables. For an arbitrary function of four variables such a representation has the form

$$f(x_1, x_2, x_3, x_4) = \sum_{r=1}^4 h^r[x_4, g_1^r(x_1, x_2, x_3), g_2^r(x_1, x_2, x_3)].$$

The question of the possibility of representing an arbitrary continuous function of three variables in the form of a superposition of continuous functions of two variables remains open. The proof of the possibility of such a representation would yield a complete solution of Hilbert's 13th problem [1] in the sense of establishing the hypothesis stated by Hilbert. Theorem 2 shows only that representation of an arbitrary continuous function of three variables in the form of a superposition of continuous functions of two variables is in fact possible if as auxiliary variables one admits variables running through a one-dimensional configuration somewhat more complicated than an interval of the real line, and in fact, the universal tree (a tree is a locally connected continuum not containing a homeomorphic image of a circle; as Menger [2] showed, there is a universal tree containing homeomorphic images of all trees).

In the sequel, k, m, n, r will denote natural numbers; $a, b, c, C, d, M, R, x, y, u, v, f, F, g, h, \epsilon, \delta, \rho$ will denote real numbers; ξ, ϕ, ψ will denote elements of trees; E^n will denote the n -dimensional unit cube, $0 \leq x_i \leq 1$; $i = 1, \dots, n$.

Theorem 1.

a) For arbitrary $n \geq 2$, there are continuous functions

$$\phi^1, \dots, \phi^{n+1}$$

defined on E^n with values in the universal tree Ξ such that every continuous real-valued function f defined on E^n has the form

$$f(x_1, \dots, x_n) = \sum_{r=1}^{n+1} h_f^r[\phi^r(x_1, \dots, x_n)],$$

where the real functions $h_f^r(\xi)$ are defined and continuous on Ξ .

b) Here the functions h_f^r can be chosen so that they will depend continuously on f in the sense of the topology of uniform convergence in the spaces of continuous functions on E^n and Ξ .

Theorem 2 follows almost immediately from Theorem 1.

Theorem 2. For arbitrary $n \geq 3$, there exist functions

$$\phi^1, \dots, \phi^n$$

defined and continuous on E^n with values in Ξ , such that every continuous function f defined on E^n is representable in the form

$$f(x_1, \dots, x_n) = \sum_{r=1}^n h^r[x_n, \phi^r(x_1, \dots, x_{n-1})],$$

where the real functions $h^r(x, \xi)$ are defined and continuous on the product $E^1 \times \Xi$.

The universal tree Ξ (see [2]) can be realized in the form of a continuum lying in the unit square E^2 . Writing g_1^r and g_2^r for the coordinates of the point ϕ^r , we obtain this assertion as an immediate consequence of Theorem 2:

Theorem 3. For arbitrary $n \geq 3$, there exist continuous real functions

$$g_1^1, \dots, g_1^n; g_2^1, \dots, g_2^n,$$

defined on E^n such that every continuous function f given on E^n is representable in the form

$$f(x_1, \dots, x_n) = \sum_{r=1}^n h^r[x_n, g_1^r(x_1, \dots, x_{n-1}), g_2^r(x_1, \dots, x_{n-1})],$$

where the functions h^r are defined and continuous on E^3 .

For $n = 3$, Theorem 3 is trivial: it has real interest only for $n \geq 4$.

It remains to sketch the method of proof of Theorem 1. The following lemma is the starting point for this.

Basic Lemma. For arbitrary $n \geq 2$, one can define on E^n a system of functions

$$u_{km}^r(x_1, \dots, x_n)$$

with indices r, k, m running through values in the intervals

$$1 \leq r \leq n+1, 1 \leq k < \infty, 1 \leq m \leq m_k,$$

that possesses the following properties:

- 1) $u_{km}^r \geq 0$;
- 2) $u_{km}^r \neq 0$ only on a set G_{km}^r of diameter d_k , where $d_k \rightarrow 0$ as $k \rightarrow \infty$;
- 3) two sets G_{km}^r and $G_{k'm'}^r$, with common indices r and k and $m' \neq m$ do not intersect;

4) for arbitrary k and at every point $P \in E^n$,

$$c \leq \sum_{r=1}^{r=n+1} \sum_{m=1}^{m=m_k} u_{km}^r \leq C,$$

where c and C do not depend upon k ;

5) the function u_{km}^r is constant on each $G_{k'm'}^r$ with the same index r for

$k' > k$ and arbitrary m' .

The construction of the system of functions u_{km}^r cannot be set down within the limits of this note. In the sequel, we shall suppose that this system of functions is given.

Lemma 1.

a) An arbitrary continuous function f defined on E^n can be represented in the form

$$f(P) = \sum_{k=1}^{\infty} \sum_{r=1}^{n+1} \sum_{m=1}^{m_k} a_{km}^r(f) u_{km}^r(P), \quad (1)$$

where the coefficients $a_{km}^r(f)$ do not depend upon P .

b) The coefficients $a_{km}^r(f)$ can be chosen in the form of continuous functions of f and indeed so that

$$|a_{km}^r(f)| \leq a(\mathfrak{F}), \quad \sum_{k=1}^{\infty} a_k(\mathfrak{F}) < \infty$$

on every family \mathfrak{F} of uniformly bounded and equicontinuous functions f .

The proof of Lemma 1 is based on properties 1), 2), and 4) of the system u_{km}^r and begins with estimates of the remainder term R in the representation

$$f(P) = \sum_{r=1}^{n+1} \sum_{m=1}^{m_k} b_m^r u_{mk}^r(P) + R,$$

where

$$b_m^r = \frac{1}{C} f(P_{km}^r),$$

and P_{km}^r are arbitrary points belonging respectively to the sets G_{km}^r . It is easy to show that for an appropriate choice of the coefficients b_m^r ,

$$|R| \leq (|1 - \frac{c}{C}| + \delta_k) M,$$

where

$$M = \sup_{P \in E^n} |f(P)|, \quad \delta_k = \sup_{\rho(P, P') \leq d_k} |f(P) - f(P')|.$$

A complete proof of Lemma 1 cannot be given here.

We now write the decomposition (1) in the form

$$f(P) = \sum_{r=1}^{n+1} f^r(P), \quad (2)$$

$$f^r(P) = \sum_{k=1}^{\infty} \sum_{m=1}^{m_k} a_{km}^r(f) u_{km}^r(P).$$

The following property of the functions f^r can be easily inferred from properties 2), 3), and 5) of the system u_{km}^r .

Lemma 2. The function $f^r(P)$ is constant on every component of an arbitrary level set of the function

$$F^r(P) = \sum_{k=1}^{\infty} \frac{1}{k^2} \sum_{m=1}^{mk} u_{km}^r(P).$$

We now remark, as was shown by A. S. Kronrod [3], that the components of level sets of an arbitrary continuous function form a tree in a certain natural topology. The tree of components of the level sets of the function F^r will be denoted by Ξ^r , and these trees Ξ^1, \dots, Ξ^{n+1} are then mapped by homeomorphisms

$$\psi_r(\Xi^r) = \Xi_0^r \subseteq \Xi$$

onto pairwise disjoint subsets of the universal tree Ξ . We set

$$\phi^r(P) = \psi_r(\xi^r), \text{ if } P \in \xi^r \in \Xi^{r0},$$

and define continuous functions $h^r(\xi)$ on Ξ such that for $\xi \in \Xi_0^r$,

$$h^r(\xi) = y, \text{ if } f^r(P) = y \text{ for } P \in \psi_r^{-1}(\xi).$$

It is easy to verify that

$$f^r(P) = h^r[\phi^r(P)]. \quad (3)$$

Formulas (2) and (3) lead us to a proof of assertion a) of Theorem 1. Assertion b) of Theorem 1 is proved on the basis of assertion b) of Lemma 1.

In conclusion we list without proof the following assertion.

Theorem 4. For arbitrary $n \geq 2$ and $\epsilon > 0$, for every function f defined and continuous on E^n , there exist polynomials

$$b(u_1, \dots, u_{n-1}), a_r(x), c_r(x); r = 1, \dots, n+1,$$

such that at all points $P \in E^n$

$$|f(P) - \tilde{f}(P)| < \epsilon,$$

where

$$\tilde{f}(x_1, \dots, x_n) = \sum_{r=1,2} a_r(x_n) b[c_r(x_n) + x_1, \dots, c_r(x_n) + x_{n-1}]. \quad (4)$$

In the case $n = 3$, upon setting

$$d(u, v) = u + v, g_r(x, y) = a_r(x)y, h_r(x, x') = c_r(x) + x',$$

we obtain from (4)

$$\tilde{f}(x_1, x_2, x_3) =$$

$$d(g_1\{x_3, b[h_1(x_3, x_1), h_1(x_3, x_2)]\}, g_2\{x_3, b[h_2(x_3, x_1), h_2(x_3, x_2)]\}). \quad (5)$$

In view of Theorem 4, every continuous function of three variables can be approximated arbitrarily by an expression of the form (5), where d, g_r, b , and h_r are polynomials of two variables. This remark also illuminates from a rather new side the circle of problems relating to Hilbert's 13th problem.

BIBLIOGRAPHY

- [1] D. Hilbert, *Gesammelte Abhandlungen*, Bd. 3, J. Springer, Berlin, 1935; no. 17.
- [2] K. Menger, *Kurventheorie*, Teubner, Leipzig-Berlin, 1932; Kap. X, §6.
- [3] A. S. Kronrod, *On functions of two variables*, *Uspehi Mat. Nauk* 5(1950), no. 1 (35), 24-134. (Russian)

Translated by:
Edwin Hewitt